

NUMERIC WILDLIFE HABITAT CLASSIFICATION OF WELLS
ISLANDS, HOOD RIVER COUNTY, OREGON

by

John L. Marshall

A RESEARCH PROJECT

Submitted to

THE OREGON DEPARTMENT OF FISH AND WILDLIFE

August, 1985

ACKNOWLEDGEMENTS

The completion of this project was made possible through the concerted effort of a group of volunteers, cooperation from the Oregon Department of Fish and Wildlife, the US Army Corps of Engineers, and the Port of Hood River.

Kathy Peck, naturalist for the Corps of Engineers, Dave Olcott, science teacher for Centennial School District, and Lucile Wyers, retired citizen of Hood River County, all contributed their time and expertise in collecting field data for the habitat classification. Jeff Barker, post graduate botanist from Washington State University, identified much of the flora collected from the islands.

John Beck and Jim Torland, wildlife biologists for the Oregon Department of Fish and Wildlife, provided boat transportation and helped collect field data for the habitat classification. John Beck also photographed the study area during a low elevation flight over the islands. The resulting photograph proved invaluable for delineating habitat boundaries during the field investigations. Nearly all wildlife observations were made by the biologists.

John Thies and Vicky Collins, from the Corps of Engineers Photogrammetry Section; Portland Office, enlarged and sent the imagery used in the study.

Finally, I would like to thank Jim O'Banion, Port of Hood River Manager, for permitting us access to the islands.

NUMERIC WILDLIFE HABITAT CLASSIFICATION OF WELLS
ISLANDS, HOOD RIVER COUNTY, OREGON

Resource managers and planners are often confronted with issues requiring decisions on the effects future management practices may have on wildlife habitat. This research provides numeric base line data on wildlife habitat characteristics (e.g., vegetation, exposed substrate, and water) of Wells Island and its two satellite islands (referred to here as Wells Islands). The data are used to construct a wildlife habitat classification. There is no attempt to evaluate the habitat for its general or species-specific wildlife value. The classification merely represents a frozen picture of existing wildlife habitat characteristics of the islands.

The objectives of this project are to:

Stratify wildlife habitat on Wells Islands using black and white air photo imagery at a scale of 1:5000.

Sample each of the units delineated on the air photo.

Use a modified numeric classification system (Marshall 1985) to classify each of the units represented by the samples.

Develop a wildlife habitat classification map based on the numeric classification employed in the unit analysis.

A multi-step process described by Marshall (1985) was modified and used to classify the wildlife habitat on the islands. Habitat boundaries were provisionally delineated on a black and white aerial photo of the islands (Corps of

Engineers-82-358). The original imagery was at a scale of 1:48,000. This imagery was subsequently enlarged to approximately 1:5,000. Since the islands were near the center of the original photo, the enlarged photo was judged to have negligible distortion and was found acceptable for mapping and classification purposes /1 . Boundaries were drawn around discernible vegetation, water, and exposed substrate units (here referred to as image units) on clear acetate overlaying the photo. Characteristics used to determine boundaries included image: density, texture, tone, and shape.

Sampling

Each image unit was sampled in the field. Data collected at each sample (Figures 1 and 3) included:

1. Unit wildlife habitat characteristics
 - A. Plant species present
 - B. Water bodies (hydrologic traits) present
 - C. Exposed substrate types present
 - D. Wildlife species observed

2. Numeric and qualitative measurement of unit characteristics
 - A. Cover class and cover class mid-points for unit characteristics A, B, and C (Table 1)
 - B. Moisture tolerance and life form for each plant species present:

/1 A low elevation air photo, taken with a 35 mm camera, was also used to help determine habitat boundaries or image units.

Species Moisture Tolerance	Life Form
obligate hydrophyte (h)	Tree /2
facultative hydrophyte (h/m)	Shrub
mesophyte (m)	Emergent

- C. Mean depth of each water body (hydrologic trait) present
- D. Height class for each plant species present (Table 1)
- E. Number of individuals of each wildlife species observed (Figure 3)
- o Time of day wildlife observations were made
 - o Sex and approximate age of animals
 - o Brief description of general behavior, i.e., feeding, incubating, defending territory, etc.
 - o Brief description of any unit characteristic(s) obviously important to the observed animal(s), i.e., trees used for perching or nesting, plants or animals used as a food source, plants used for cover, etc.

Table 1. Height and Cover Class Parameters (Kuchler 1966).

Height	Height Class	% Cover	Cover Class	(% Mid-Point)
35-45m	8	75-100	5	88
20-35m	7	50-75	4	63
10-20m	6	25-50	3	38
5-10m	5	5-25	2	15
2- 5m	4	0- 5	1	3
.5- 2m	3			
.1-.5m	2			
< .1m	1			

/2 Trees constitute predominantly woody plants greater than 2 meters in height.

All tree height measurements were determined using a clinometer. Cover class measurements were subjective estimates made during a visual surveillance of an undefined area within each image unit boundary.

One sample was taken for each image unit. Thirty image units were delineated on the photo. Each unit was photographed in the field with a 35 mm camera.

Importance Index

After the data were compiled for each sample, an importance index was calculated and used to organize the data and classify the wildlife habitat on the islands. The classification is based on structural characteristics of vegetation, depth and area of water bodies (hydrologic traits), and area of substrate. /3 The importance index is determined by running the data from each sample through six steps:

1. Group plant species by life form and moisture tolerance
2. Sum percent cover class mid-points for each life form, hydrologic trait, and substrate type.

/3 The hydrologic traits "Shallow" and "Deep Open Water" and all substrate types were arbitrarily assigned a significance coefficient of 4. Significance coefficients for hydrologic traits and substrate types serve the same function as height class numbers for vegetation life forms. That is, they weight the importance value of their respective habitat characteristics. Relatively high significance coefficients were selected on the premise that hydrologic and substrate regimes heavily influence the physical and biological character of the units they occupy. Subsequently, they are important influences on wildlife associated with those units. Therefore, the importance index reflects, albeit crudely, ecological factors that contribute to the character of the unit.

3. Determine the mean height class number for each life form and the significance coefficient for each hydrologic trait and substrate type. This is the height class/significance coefficient index.
4. Determine an importance value for each life form by multiplying the sum of the life form cover class mid-points by its height class index. The importance value for hydrologic traits and substrate types are derived by multiplying their respective cover class mid-points by their corresponding significance coefficients.
5. Determine sum of importance values for all life forms, hydrologic traits, and substrate types.
6. Determine a relative importance value for each life form, hydrological trait, and substrate type by dividing each respective importance value by the sum of all importance values. This is the importance index number.

The process by which importance indexes were derived can be followed on Figure 1. Data from sample 23, image unit 19 (units were not sampled in the order of their assigned numbers) are displayed to allow the reader to follow the procedure. This form was used for each image unit sample.

Wildlife Habitat Classification

The subsequent task of defining the relative importance of various wildlife habitat classes in each sample was accomplished using Figure 2. The importance index derived for each wildlife habitat component on Figure 1 was transferred to the corresponding life form, hydrologic trait, and/or substrate type on Figure 2. A "dominance threshold index", within the range of 0 to 1, was arbitrarily selected to categorize the dominant habitat class. Any life form, hydrologic trait, or substrate type that obtained an importance index greater than or equal to the threshold was considered

a dominant habitat class. The threshold was 0.38. A "subordinate threshold index" of 0.14 was selected to delineate subdominant classes. While definitions of threshold indexes were arbitrary, they were based on many hours of field observation and working with the data.

Using the data in unit 19 (Figures 1 and 2), the Moist Forest wildlife habitat characteristic has an importance index of .56; qualifying it for dominant class status. The Exposed Substrate (3) habitat characteristic (sand) received an importance index of .37; qualifying it as a subordinate wildlife habitat class (subclass) in the unit.

An upper case letter code (A-N) was used to define habitat classes in each unit. Habitat classes were represented fractionally in the code with increasing importance to the left. A slash separates the dominant classes from the subordinate. A dash separates codominant or cosubordinate classes. The code for unit 19 is:

A / X K

K A - Moist Forest (dominant)
X - Sand (subordinate)

Following Larson (1976) and Cowardin et al. (1979), each class is modified by descriptive components of the chief habitat characteristics of the unit, i.e., short narrow leaf emergents, floating vascular emergents, dead trees, etc. Modifiers are represented here by a lower case letter code and are listed in order of decreasing importance to the

Physiographic Province: Columbia River Gorge	Land Form: Island	Area Type: Upland nonwetland Upland wetland Bottomland nonwetland Bottomland wetland	Wetland System/Subsystem: Marine / Subtidal, Intertidal Estuarine / Subtidal, Intertidal Riverine / Tidal, Lower Perennial, Upper Perennial, Intertidal. Lacustrine / Limnetic, Littoral Palustrine						
Location: Columbia River (Bonneville Pool) River Mile 148			Island: Wells I	Sampler: Z3	Unit: 19	Date: 6-28-85			
(Step 1)									
Plant Species /1	Plant Moisture Tolerance h/h/a/a	Hydro-logic Trait (N.T.)	Exposed Substrate Type (S.T.)	Site Characteristics				Exposed Substrate	
				Cover Class and Cover Class Mid-Points (Percent)					
				Life Form (L.F.)		Hydrologic Trait (H.T.)			
				Trees (T)	Shrubs (S)	Herbs (H)	Open Water Wetland		
							Shallow (< 2 m DM1)	Deep (> 2 m DM2)	
SALIX	X		5 RB						Sr CIG S Si Cl
SOOC	X				2	15			
POTR	X			1	3				
COST	X			1	3				
			SAND						5
				88	6	15			5
(Step 2) Sum the percent cover mid-points for each site characteristic (L.F., H.T., and S.T.).									

(Step 3) HEIGHT CLASS/SIG. COEFF. INDEX
Find the mean H.C. for each respective L.F. and the significance coefficient for each S.T. and H.T.

* 5=88

Plant Species	Hydrologic Trait	Substrate Type	Life Form Height Class			Hydrologic Trait Sig. Coeff. (S.C.)	Exposed Substrate Type Sig. Coeff. (S.C.)							H.C./S.C. Index /2																											
			T	S	H		Sh	C	G	S	Si	Cl	T	S	H	DM1	DM2	SE	C	G	S	Si	Cl																		
SALIX			6																																						
SOOC				3																																					
POTR				3																																					
COST				3																																					
		SAND																																							

(Step 4) IMPORTANCE VALUE
For each L.F., H.T., and S.T., multiply the mean of cover class mid-points by their respective H.C. and S.C. indexes.

(Step 5)
Sum the importance values.

(Step 6) IMPORTANCE INDEX
Divide each importance value by the sum of importance values.

Trees	Shrubs	Herbs	Open Water		Exposed Substrate						
			Shallow (< 2 m)	Deep (> 2 m)	Sh	C	G	S	Si	Cl	
328	18	45								352	
943.00											
.56	.02	.05								.37	

/1 Key to plant acronym is in Appendix 2.

/2 The deep and open water hydrologic traits were arbitrarily assigned the numerical significance coefficient of 4. This was done on the grounds that these traits significantly affect the physical and biological character of wetlands. The exposed substrate types were also assigned a common significance coefficient of 4. This was done to reflect the effect exposed substrate has on the physical and biological character of both wetlands and uplands.

Figure 1. Importance indexes for Wells Island unit number 19.

Physiographic Province: Columbia River Gorge		Land Form: Wells Island No. 1		Area Type: bottomland wetland		Wetland System/Subsystem: Lacustrine/Littoral	
Location: Columbia River (Bonneville Pool) River Mile 168				Water Regime: 7		Special Modifier: 2	
Date: 06-28-85		Unit: 19		Samples: 23		Code: Abd/Kh	
A	B	C	D	E	F	G	
T (1) 60% h) > 2 a Moist Forest	T (1) 60% a) > 2 a Mesic Forest	S (1) 60% h) < 2 a Moist Shrub- land	S (1) 60% a) < 2 a Mesic Shrub- land	N (1) 60% h) Frequent Emergent Wetland	N (1) 60% h/a) Infrequent Emergent Wetland	N (1) 60% a) Meadow or Mesic Herb	
.56		.02			.05		
a. Broad Leaf Decid. b. Narrow Leaf Decid. I c. Evergr. d. Dead I	a. Broad Leaf Decid. b. Narrow Leaf Decid. c. Evergr. d. Dead	a. Broad Leaf Decid. I b. Narrow Leaf Decid. c. Evergr. d. Dead e. Lianas	a. Broad Leaf Decid. b. Narrow Leaf Decid. c. Evergr. d. Dead e. Lianas	a. Short Narrow Leaf < 2 a b. Tall Narrow Leaf > 2 a c. Short Broad Leaf < 2 a d. Tall Broad Leaf > 2 a e. Robust f. Persist. g. Nonpers. h. Floating Vasc.	a. Short Narrow Leaf < 2a I b. Tall Narrow Leaf > 2a c. Short Broad Leaf < 2 a d. Tall Broad Leaf > 2 a e. Robust f. Persist. g. Nonpers.	a. Short Narrow Leaf < 2a b. Tall Narrow Leaf > 2a c. Short Broad Leaf < 2 a d. Tall Broad Leaf > 2 a e. Robust f. Persist. g. Nonpers.	
H	I	J	K	L	M	N	O
ES(1) > 50% Stone	ES(2) > 50% Cobble	ES(3) > 50% Gravel	ES(4) > 50% Sand	ES(5) > 50% Silt	ES(6) > 50% Clay	OW(1) Shallow Open Water Wetland < 2 meters	OW(2) Deep Open Water Wetland > 2 meters
			.37				
a. Consol. b. Fragments c. 15-50% d. Cobble e. Gravel f. Sand g. Silt h. Clay i. Very Wet j. Mod Wet k. Dry	15-50% a. Stone b. Gravel c. Sand d. Silt e. Clay f. Very Wet g. Mod. Wet h. Dry	15-50% a. Stone b. Cobble c. Sand d. Silt e. Clay f. Very Wet g. Mod. Wet h. Dry	15-50% a. Stone b. Cobble c. Gravel d. Silt e. Clay f. Very Wet g. Mod. Wet h. Dry I	15-50% a. Stone b. Cobble c. Gravel d. Sand e. Clay f. Very Wet g. Mod. Wet h. Dry	15-50% a. Stone b. Cobble c. Gravel d. Sand e. Silt f. Very Wet g. Mod. Wet h. Dry	a. Algal b. Aquatic Hess c. Sub- merged Vascular	a. Algal b. Aquatic Hess c. Sub- merged Vascular

Figure 2. Wildlife habitat classification of Wells Island unit number 19.

Wildlife Species Observed	Number of Individuals	Sex/Age	Time of Day	Brief Description of Behavior	Brief Description of Site Characteristics Obviously Important to Animal(s)
Canada Goose	3 eggs	N/A	4 pm	N/A	Exposed sand levee Cover & Water
Black-Capped Chickadee	2	Adult Male & Female	4pm	Diversion act	Dead willow snags for cavity nesting

Figure 3. Wildlife observations.

right of the class or subclass they describe. /4

They are listed for unit 19 under the following letter code:

Ad[6]e[2]ac / Ja[6]b[2]

A = Moist Forest
d[6] = water regime [Intermittently Flooded]
e[2] = special modifier [Impounded]
a = deciduous trees
c = dead trees

X K = Sand
a[6] = water regime [Intermittently Flooded]
b[2] = special modifier [Impounded]

The modifiers in the Cowardin (1979) classification apply specifically to wetlands. While wetland wildlife habitats are recognized as being associated with the islands, the modifiers here are applied to both nonwetland and wetland units. All wetlands associated with the islands are considered "Lacustrine" (subsystem "Littoral") or "Palustrine" wetlands under the Cowardin et al. (1979) classification (Appendix 1).

The water regime modifier consists of eight descriptors. /5 All but the eighth descriptor were adopted from Cowardin et al. (1979):

1. **Permanently Flooded**

Water that covers the land surface throughout the year in

/4 While the "water regime" modifier and the "special modifier" are always listed first and second in the code respectively, they are not considered in the hierarchy of relative importance as are all other modifiers. Their position does, however, represent the major role water regime and water management play in determining the character of the unit.

/5 These descriptors were called modifiers in Cowardin et al. (1979).

all years. Vegetation is composed of obligate hydrophytes.

2. **Semipermanently Flooded**

Surface water persists throughout the growing season in most years. When surface water is absent, the water table is usually at or very near the land surface.

3. **Seasonally Flooded**

Surface water is present for extended periods early in the growing season, but is absent by the end of the season in most years. When surface water is absent, the water table is often near the land surface.

4. **Saturated**

The substrate is saturated to the surface for extended periods during the growing season, but surface water is seldom present.

5. **Temporarily Flooded**

Surface Water is present for brief periods during the growing season, but the water table usually lies well below the soil surface for most of the season. Plants that grow both in uplands and wetlands are characteristic of the temporarily flooded regime.

6. **Intermittently Flooded**

The substrate is usually exposed, but surface water is present for variable periods without detectable seasonal periodicity. Weeks, months, or even years may intervene between periods of inundation. The dominant plant communities under this regime may change as soil moisture conditions change. Some areas exhibiting this regime do not fall within the Cowardin (1979) definition of wetland because they do not have hydric soils or support hydrophytes.

7. **Artificially Flooded**

The amount and duration of flooding is controlled by means of pumps or siphons in combination with dikes or dams. The vegetation growing on these areas cannot be considered a reliable indicator of water regime. Examples of artificially flooded wetlands are some agricultural lands managed under a rice-soybean rotation, and wildlife management areas where forests, crops, or pioneer plants

may be flooded or dewatered to attract wetland wildlife. Neither wetlands within or resulting from leakage from man-made impoundments, nor irrigated pasture lands supplied by diversion ditches or artesian wells are included in this modifier.

8. Permanently Exposed

Upland areas that are outside of the 100 year flood fringe and have moderate to well drained soils. Dominant plant communities are mesic or xeric site indicators. Herbaceous plant communities indicate flood regime on sandy soils.

In addition to the above descriptors, Cowardin et al. (1979) lists six special modifiers. These modifiers are also recognized in this classification:

1. Excavated

Lies within a basin or channel excavated by man.

2. Impounded

Created or modified by a man-made barrier or dam which purposefully or unintentionally obstructs the outflow of water. Both man-made dams and beaver dams are included.

3. Diked

Created or modified by a man-made barrier or dike designed to obstruct the inflow of water.

4. Partly Drained

The water level has been artificially lowered, but the area is still classified as a wetland because soil moisture is sufficient to support hydrophytes. Drained areas are not considered wetlands if they can no longer support hydrophytes.

5. Farmed

The soil surface has been mechanically or physically altered for production of crops, but hydrophytes will become reestablished if farming is discontinued.

6. Artificial

Refers to substrates classified as Rock Bottom, Unconsolidated Bottom, Rocky Shore, and Unconsolidated Shore that were emplaced by man, using either natural materials such as dredge spoil or synthetic materials such as discarded automobiles, tires, or concrete. Jetties and breakwaters are examples of Artificial Rocky Shores. Man-made reefs are an example of Artificial Rock Bottoms.

The classification of each unit (without modifiers and descriptors) is recorded in Table 2. Figure 4 shows the distribution of units sampled on Wells Islands. Figure

Table 2. Wells Islands Wildlife Habitat Unit Classification.

Unit	Class	Subclass
1	A	G- X K
2	M	A
3a	G	A /1
3b	A	E
4	G	
5	G	
6	G	X K
7	A	G
8	G-D	
9	A	G
10	X K	G
11	G	
12	A	F
13	A	F-C
14 /2		
15	G	
16	A	G
17	X K	
18	A	G

/1 Units 3a and 3b were split in the field due to the obvious differences in habitat characteristics.

/2 Unit 14 was joined with unit 15 in the field due to the obvious similarities in habitat characteristics and their spatial proximity.

Table 2. (Continued) Wells Islands Wildlife Habitat Unit Classification.

Unit	Class	Subclass
19	A	* K
20 /3		
21	A	* K
22	A	G
23a	A	G
23b	* K	A
24	A	G
25	G	D
26	U /4	
27	A	G
28	A	G
29	B	G
30	G	

/3 Unit 20 was joined with number unit 11 in the field.

/4 Unit 26 has not been classified.

5 shows the distribution of dominant and subordinate wildlife habitat classes for the islands.




Figure 4. Wells Islands image units.




Figure 5. Wells Islands wildlife habitat classification.

LITERATURE CITED

- Corps of Engineers. 1982. Air photo, 82-358.
- Cowardin, L.M., U. Carter, F. Golet, and E. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States, Biological Services Program, U.S. Fish and Wildlife Serv. Washington, D.C., 103 p.
- Donahue, R.L., L. Robertson, and J. Schickluna. 1971. Soils: an introduction to soils and plant growth, Prentice Hall, Inc., Englewood Cliffs, New Jersey.
- Kuchler, A.W. 1966. Analyzing the physiognomy and structure of vegetation, Ann. Assoc. Amer. Geog., 56:112-126.
- Larson, J.S. 1976. Models for assessment of freshwater wetlands, Water Resource Research Center, Univ. of Mass. Pub. No. 32., 86 p.
- Marshall, J.L. 1985. Value assessment of Jackson-Frazier Wetland, Benton County, Oregon: a case study, unpublished thesis, Dept. of Geography, Oregon State Univ., 147 p.